

***Chandra* Detection of Local O VII He α Absorption Along the Sight Line Toward 3C 273**

Taotao Fang^{1,2}, Kenneth R. Sembach³, AND Claude R. Canizares¹

ABSTRACT

With the *Chandra* X-ray Telescope we have detected a zero-redshift O VII He α absorption line along the sight line toward 3C 273. This line detection is highly significant, with a signal-to-noise ratio (S/N) of 6.4. We explore two models, which associate this line with (1) the intragroup gas in the Local Group, and (2) the hot halo gas in the vicinity of our Milky Way. In the first model, we find that for a standard β -model of the gas distribution in the Local Group, the temperature is constrained to $2.3 \times 10^5 < T < 1.2 \times 10^6$ K and the baryon overdensity is $\delta_b \sim 100$; both results are consistent with the properties of the so-called warm-hot intergalactic medium (WHIM) predicted by cosmological simulations. We also find the core radius of the gas distribution should be > 100 kpc. In the second model we discuss several possible Galactic origins for the absorption, and we comment on the possibility that the O VII is associated with the O VI absorption observed in this direction by the Far Ultraviolet Spectroscopic Explorer (*FUSE*). We find that there is a strong indication that collisional ionization is the dominant ionization source for the observed absorption.

Subject headings: ISM: cloud — Local Group — intergalactic medium — quasars: absorption lines — quasars: individual (3C 273) — X-rays: individual (3C 273)

1. Introduction

Highly ionized resonance absorption lines from heavy elements (such as C IV, Si IV, N V, O VI) at low redshift have been detected along many lines of sight by the Far Ultraviolet Spectroscopic Explorer (*FUSE*) and the Hubble Space Telescope (*HST*) (e.g., Tripp, Savage, & Jenkins 2000; Sembach et al. 2000; Savage et al. 2000). These species arise in

¹Department of Physics and Center for Space Research, MIT, 77 Mass. Ave., Cambridge, MA 02139,

²Department of Physics, Carnegie Mellon University, 5000 Forbes Ave., Pittsburgh, PA 15213

³Space Telescope Science Institute, 3700 San Martin Dr., Baltimore, MD 21218

the interstellar medium of the Milky Way and the low redshift intergalactic medium (IGM). Among these ions, O VI is of particular interest because it is an excellent tracer of gas at $T \geq 10^5$ K. The detection of O VI in many directions (Savage et al. 2002; Sembach et al. 2002; Wakker et al. 2002) indicates the existence of hot gas in the halo of the Milky Way, in high velocity clouds surrounding the Milky Way, or even in the Local Group (Nicastro et al. 2002).

Recently, with the unprecedented high resolution of the *Chandra* X-ray Telescope, Nicastro et al. (2002) and Fang et al. (2002) reported the first detection of X-ray absorption lines from higher ionization species, O VII and O VIII, at $z \approx 0$ along the sight line toward the quasar PKS 2155-304. Mathur, Weinberg, & Chen (2002) reported possible detection of such X-ray absorption feature in the spectrum of H 1821+643. Kaspi et al. (2002) also reported the detection of zero-redshift O VII in a long exposure (~ 900 ksec) of the bright Seyfert galaxy NGC 3783 with *Chandra*. Both O VII and O VIII trace gases with temperatures $\gtrsim 10^6$ K. Therefore, it is of great interest to see whether the higher temperature gas is related to the gas that produces the UV absorption lines and whether there is a multi-phase distribution of hot gas in the vicinity of the Milky Way. It is also possible that the X-ray absorption is associated with hot intragroup gas in the Local Group. If so, this might provide important information about the “missing baryons” at low redshift. Cosmological simulations indicate that a large fraction of neutral gas at high redshift is heated to high temperatures ($10^5 < T < 10^7$ K) at $z \sim 0$ (Cen et al. 1995; Ostriker & Cen 1996; Fukugita, Hogan, & Peebles 1998; Cen & Ostriker 1999); gas in the higher end of this temperature range has so far escaped detection at low redshift. Simulations predict that these “missing baryons” are distributed in intergalactic space in the form of diffused filamentary structures (the warm-hot intergalactic medium, or WHIM) that connect virialized clusters and/or groups of galaxies with overdensities of $\delta_b \sim 5 - 200$.

In this *Letter* we report the detection of a $z \sim 0$ O VII He α resonance absorption line in the *Chandra* spectra of 3C 273. We explore two possible sources of this absorption: (1) intragroup gas in the Local Group, and (2) hot gas in the Milky Way halo and interstellar medium along the sight line.

2. Data Analysis

3C 273 ($l = 289.95^\circ$, $b = 64.36^\circ$, $z_{QSO} = 0.158$) is one of the brightest quasars in the sky at X-ray wavelengths and has been repeatedly observed as a *Chandra* calibration target. It was observed with the *Chandra* Low Energy Transmission Grating Spectrometer (LETGS) and the Advanced CCD Imaging Spectrometer (ACIS) on 31 October 2000 (obs.

#1198), 13 June 2001 (obs #2464), and 15 June 2001 (obs #2471), with an exposure time of 40, 30 and 30 ksec, respectively. See the *Chandra* Proposers’ Observatory Guide at <http://asc.harvard.edu/> for more information about *Chandra* and its instrumentation. In the analysis and discussions that follow, we adopt $H_0 = 70 h_{70} \text{ km s}^{-1} \text{ Mpc}^{-1}$. Unless otherwise stated, errors are quoted at 90% confidence.

We reduced the Level 1 data using standard CIAO tools⁴, and restricted all spectra to the 2–42 Å wavelength range to avoid complexities at longer wavelengths from the instrumental Carbon K edge. We summed the events in 0.025 Å bins, which are half the width of the instrumental line response function (FWHM ~ 0.05 Å across the bandpass). To remove the underlying broad-band continuum behavior for each dataset, we divided each spectrum by the best fit power-law, with a correction for absorption by the foreground Galactic interstellar medium. We then fit the remaining residuals with a sixth-order polynomial. These procedures remove spectral features larger than ~ 8 Å in width (e.g., broad 5–10% calibration uncertainties) but preserve narrow line features. We combined the data from the three observations as a weighted sum accounting for differences in exposure times and effective areas.

All of the continua are well described by a single power law with a similar power-law index. The power-law photon indices are $\Gamma_{1198} \sim 2.08 \pm 0.36$, $\Gamma_{2464} \sim 1.86 \pm 0.40$, and $\Gamma_{2471} \sim 1.82 \pm 0.48$, assuming absorption by a Galactic neutral hydrogen column of $N_{HI} = 9 \times 10^{19} \text{ cm}^{-2}$ (Lockman & Savage 1995). The October 2000 observation exhibits the strongest soft X-ray flux; $F_{0.1-2.4 \text{ keV}} = (9.3, 7.9, 7.8) \times 10^{-11} \text{ ergs cm}^{-2} \text{ s}^{-1}$ for observations #1198, #2464, and #2471, respectively. The total counts per bin in the continuum decline smoothly from ~ 150 at 18 Å to ~ 40 at 23 Å, with ~ 70 at 21.6 Å.

After a blind search for any statistically significant absorption features in the 2–42 Å region of the LETGS spectral bandpass (Figure 1), we identified an absorption feature with S/N ~ 6.4 at 21.60 ± 0.01 Å, which is almost identical to the rest wavelength of O VII He α (see Table 1). We fit this feature with single-component Gaussian models using ISIS⁵, taking into account the instrumental line response function (Table 1). We also examined other *Chandra* LETGS archival data for 3C 273 and found a similar feature at the same wavelength with a different focal plane detector - the High Resolution Camera (HRC).

⁴Chandra Interactive Analysis of Observations (CIAO)”, see <http://asc.harvard.edu/ciao/>

⁵Interactive Spectral Interpretation System, see Houck & Denicola 2000.

3. Local O VII He α Absorption

Based on the detected line equivalent width (W_λ), we can estimate the implied column density of O VII (Spitzer 1978). Since the line is unresolved, the Doppler b -parameter must be $b = \sqrt{2}\sigma < 400 \text{ km s}^{-1}$ (90% confidence). If $b > 200 \text{ km s}^{-1}$, then the line is unsaturated and $N(\text{O VII}) = 1.8_{-0.7}^{+2.7} \times 10^{16} \text{ cm}^{-2}$. The line becomes saturated at $b \sim 100 \text{ km s}^{-1}$, giving $N(\text{O VII}) \sim 10^{17} \text{ cm}^{-2}$. At $b < 100 \text{ km s}^{-1}$ the line is heavily saturated, but this can be ruled out by the lack of higher Lyman series lines in our spectrum. From the non-detection of local O VII He β line at $18.6288 \text{ \AA}$, we estimate a 4σ equivalent width upper limit of $11.9 \text{ m\AA}$. Based on a curve-of-growth calculation for O VII He β , we find that even in the case for which the line is heavily saturated ($b < 100 \text{ km s}^{-1}$), the He β upper limit restricts the O VII column density to $< 2 \times 10^{16} \text{ cm}^{-2}$. We therefore adopt $N(\text{O VII}) = 1.8_{-0.7}^{+0.2} \times 10^{16} \text{ cm}^{-2}$ in the following discussion.

Highly ionized metals such as C IV and O VI at $z \approx 0$ have been detected repeatedly through their absorption-line signatures along many lines of sight (see Savage et al. 2002, Sembach et al. 2002, and references therein). Unlike these lower ionization species, which typically trace transition temperature gas with $T \sim (1 - 5) \times 10^5 \text{ K}$, the O VII detected with *Chandra* indicates the existence of the gas with temperatures $T \sim 10^6 \text{ K}$. This high temperature gas traced by O VII is unlikely to be produced in the nearby interstellar medium (ISM), although we cannot rule out the possibility of a nearby supernova remnant origin. It is more plausible that the He-like Oxygen is produced in hot Galactic halo gas, or even in the Local Group. It is to these two possibilities that the rest of this *Letter* is devoted.

3.1. Comments on a Possible Local Group Origin

The detection of high column density O VII implies the existence of a large amount of hot gas with a temperature around 10^6 K . In the case of pure collisional ionization, temperature is the only parameter of importance over a wide range of density so long as the gas is optically thin. The O VII ionization fraction peaks at nearly 1, and exceeds 0.5, for temperatures $T \sim (0.5 - 1.3) \times 10^6 \text{ K}$ (Sutherland & Dopita 1993). However, the IGM is also photoionized by the background UV/X-ray radiation, which increases the ionization fraction at low temperatures and densities. We explored these regimes using a grid of CLOUDY models (Ferland et al. 1998) for $10^2 < T < 10^9 \text{ K}$ and $10^{-8} < n(\text{cm}^{-3}) < 10^{-2}$ with $0.2Z_\odot$. We used the UV background from Shull et al. (1999), and a power law with an exponential cutoff at 50 keV and $\alpha = 0.29$ in the X-ray spectral region (see, e.g. Fabian & Barcons 1992). We combined the ionizing UV and X-ray spectra using normalizations of $J_{13.6\text{eV}} = 1.8 \times 10^{-23} \text{ ergs cm}^{-2} \text{ s}^{-1} \text{ Hz}^{-1} \text{ sr}^{-1}$ and $J_{1\text{keV}} = 10 \text{ ph cm}^{-2} \text{ s}^{-1} \text{ keV}^{-1} \text{ sr}^{-1}$.

CLOUDY calculations indicate that the ionization fraction versus T relation begins to deviate appreciably from the pure collisional case for low density regions. At a baryon density $n_b \leq 10^{-4} \text{ cm}^{-3}$, photoionized low temperature gas ($4 \leq \log T \leq 5.5$) can produce significant amount of O VII. We also considered several other background radiation models, such as the X-ray background from Miyaji et al. (1998) and the UV background from Haardt & Madau (1996), but as shown by Chen et al. (2002) we found that the resulting differences in the predicted O VII ionization fraction are negligible.

The upper limit on the line width of $\sigma \sim 400 \text{ km s}^{-1}$ sets an maximum to the path length of $\sim 5 h_{70}^{-1} \text{ Mpc}$, since otherwise the differential Hubble flow would excessively broaden the line. We can further constrain the density of the absorbing gas, assuming it is associated with the Local Group. The path length can be set equal to the distance to the boundary of the Local Group, where the gas begins to participate in the Hubble flow. Assuming a simple spherical geometry for the Local Group, the path length is $\sim 1 \text{ Mpc}$ (see the following text). Adopting the 90% lower limit of the O VII column density, this gives $n_b > (2.2 \times 10^{-5} \text{ cm}^{-3}) Z_{0.2}^{-1} f_1^{-1} l_1^{-1}$, where $Z_{0.2}$ is the metallicity in units of 0.2 solar abundance, f_1 is the ionization fraction, and l_1 is the path length in units of 1 Mpc. This implies a baryon overdensity of $\delta_b \sim 100$, which is roughly consistent with the overdensity of the WHIM gas predicted from cosmological simulations (Cen et al. 1995; Ostriker & Cen 1996; Cen & Ostriker 1999; Davé et al. 2001).

A model characterizing the distribution of the Local Group gas is given by the standard β -model (Cavaliere & Fusco-Femiano 1976):

$$n(r) = n_0 \left[1 + \left(\frac{r}{r_c} \right)^2 \right]^{-\frac{3}{2}\beta} \quad (1)$$

where n_0 is the central gas number density, r_c is the core radius, β is the ratio of the specific kinetic energy to thermal energy used to characterize the depth of the potential well. Poor groups like the Local Group typically have shallow potentials with $\beta \sim 0.5$ (Mulchaey et al. 1996), so we adopt this value. We adopt a simplified geometry model of the Local Group, where the barycenter is located along the line connecting M31 and the Milky Way in the direction towards $(l, b) \approx (121.7^\circ, -21.3^\circ)$ at a distance of about 450 kpc from the Milky Way (Rasmussen & Pedersen 2001). At about $r \sim 1200 \text{ kpc}$, the gravitational contraction of the Local Group starts to dominate the Hubble flow, so this is typically defined as the boundary of the Local Group (Courteau & van den Bergh 1999).

Based on this simple model, we estimate the column density of O VII along the 3C 273 sight line through the Local Group by integrating the O VII number density $n_{\text{O VII}}$ out to the boundary. Figure 2 shows the O VII column density ($N_{\text{O VII}}$) versus baryon density at the

center of the Local Group (n_0). By applying a conservative upper limit of the Local Group gravitational mass and arguing that the cooling timescale should be longer than half the Hubble time, Rasmussen & Pedersen (2001) obtained an upper limit of the central density $n_0 \leq 5 \times 10^{-4} \text{ cm}^{-3}$. From Figure 2 we find a tight upper limit of temperature $T \leq 1.2 \times 10^6 \text{ K}$; at temperatures higher than $1.2 \times 10^6 \text{ K}$, the O VII ionization fraction drops quickly and O VIII starts to dominate. We also find that the temperature of the Local Group should be higher than $2.3 \times 10^5 \text{ K}$. To satisfy the observed O VII column density, the gas distribution should have a rather flat core with $r_c \geq 100 \text{ kpc}$.

3.2. Comments on a Hot Halo Gas Origin

Strong O VI absorption along the sight line towards 3C 273 was detected with *FUSE* between -100 and +100 km s^{-1} (Sembach et al. 2001). This absorption probably occurs in the interstellar medium of the Milky Way disk and halo. The column density, $\log N(\text{O VI}) = 14.73 \pm 0.04$, is the second highest O VI column recorded along any sight line through the Milky Way halo (see Savage et al. 2002). Absorption features of lower ionization species are also present at these velocities. Most notably, the C IV $\lambda\lambda 1548.195, 1550.770$ and N V $\lambda\lambda 1238.821, 1242.804$ lines are strong and track the O VI absorption closely. Between -100 and +100 km s^{-1} , $N(\text{O VI})/N(\text{C IV})$ and $N(\text{O VI})/N(\text{N V})$ are nearly constant, indicating that the species arise in the same gas along the sight line.

O VI absorption is also detected between +100 and +240 km s^{-1} in the form of a broad, shallow absorption wing extending redward of the primary Galactic absorption feature. This O VI absorption wing has $\log N(\text{O VI}) = 13.71$, or $\sim 10\%$ that of the primary absorption feature at lower velocities. This wing is not observed in other species observable with either *HST* or *FUSE*. The O VI absorption wing has been attributed to hot gas flowing out of the Galactic disk as part of a “Galactic chimney” or “fountain” in the Loop IV and North Polar Spur regions of the sky. Alternatively, the wing might be remnant tidal debris from interactions of the Milky Way and smaller Local Group galaxies (Sembach et al. 2001). Similar O VI absorption wings are seen in other directions in this general region of the sky (Sembach et al. 2002).

It is possible to associate the O VII absorption detected by *Chandra* with these highly ionized metals detected by *FUSE* and *HST*. Here, we briefly discuss several scenarios:

(1). *The O VII is related to the primary O VI feature.* In this case, $\log N(\text{O VI}) = 14.73$, and $\log [N(\text{O VII})/N(\text{O VI})] \sim 1.5$, assuming $N(\text{O VII}) = 1.8 \times 10^{16} \text{ cm}^{-2}$. This is within about a factor of 2 of the O VII/O VI ratio observed for the PKS 2155-304 absorber at $z \approx 0$ (Nicastro

et al. 2002; Fang et al. 2002) and is consistent with the idea that the gas is radiatively cooling from a high temperature (Heckman et al. 2002). This possibility is appealing since the centroids of the O VI and O VII absorption features are similar ($\sim 6 \pm 10 \text{ km s}^{-1}$ versus $-26 \pm 140 \text{ km s}^{-1}$) and the width of the resolved O VI line (FWHM $\sim 100 \text{ km s}^{-1}$) is consistent with a broad O VII feature.

However, this possibility also has drawbacks. First, the amount of O VIII predicted by the Heckman et al. (2002) cooling gas model would be ~ 3 times the observed O VII column density. However, we did not detect any O VIII Ly α absorption line at the corresponded wavelength in the *Chandra* spectrum. We estimate that the 4σ upper limit on the O VIII Ly α equivalent width is $\sim 14.20 \text{ m\AA}$, or $\sim 10^{16} \text{ cm}^{-2}$ if the line is unsaturated. Second, the amount of C IV predicted is a factor of ~ 10 times less than the O VI column density, which is a factor of 5 lower than observed. It is possible that the C IV column could be increased by including photoionization from the cooling flow itself (Benjamin & Shapiro 2000) to bring this ratio closer to the observed value. The predicted N(O VI)/N(N V) ratio of ~ 20 is about a factor of 3 higher than observed. If there are turbulent mixing layers (TMLs) occurring between the hot gas and cooler ISM material (a likely possibility), then the discrepancies in the O VI/C IV ratio might be alleviated since the standard TML models described by (Slavin, Shull, & Begelman 1993) predict considerably more C IV (but not more N V) than O VI.

(2). *The O VII is related to the high velocity O VI wing.* In this case, the O VII is associated only with the O VI absorption “wing” and not with the main O VI absorption at lower velocities. Although perhaps less likely than case (1), this may be plausible since the average velocity of the O VI absorption wing ($\sim 150 \text{ km s}^{-1}$) is still in reasonable agreement with the O VII centroid. In such a situation, $\log [N(\text{O VII})/N(\text{O VI})] \sim 2.5$. In collisional ionization equilibrium, this would imply a temperature of $> 10^6 \text{ K}$ (Shapiro & Moore 1976; Sutherland & Dopita 1993). In Figure 3 we show the 90% limit on N(O VII)/N(O VI) (the blue region), which gives a tight constraint on the temperature of the absorber: $6.0 < \log T < 6.9$. A further constraint can be achieved by considering the non-detection of O VIII Ly α absorption. The 4σ upper limit of column density requires $T \lesssim 10^{6.3} \text{ K}$; both N(O VII)/N(O VI) and N(O VIII)/N(O VII) give about the same constraint.

Another possibility is that the O VII is not related the O VI. In this case, the temperature must be high enough ($> 10^6 \text{ K}$) to prevent the production of O VI. The upper limit of the temperature can still be obtained by the N(O VIII)/N(O VII), which gives $T < 10^{6.3} \text{ K}$. This case is similar to that described in situation (2).

3.3. Comments on Other Locations

Can other sources contribute to the observed O VII column density? The Local Bubble, a large volume of hot, rarefied gas around the Sun, could be a potential source. The hot Local Bubble gas has a temperature of $\sim 10^6$ K (Snowden et al. 1998), and contains highly ionized metal-line species (see Sanders et al. 1998). However, given the rather small size of the Local Bubble ($\lesssim 250$ pc), even if the electron density is high (~ 0.005 cm $^{-3}$; Sfeir et al. 1999) and assuming all oxygen ions are in O VII, the overall contribution to the O VII column density is $\lesssim 3 \times 10^{15}$ cm $^{-2}$, or $\lesssim 17\%$ of the total O VII column density. Another potential source is the hot gas in the Galactic thick disk / low halo observed in O VI absorption (Savage et al. 2002). Given the high latitude ($b = 64.36^\circ$) of 3C 273, the path length through a scale height of this gas is ~ 2 kpc. Assuming a density of ~ 0.001 cm $^{-3}$, solar abundances, and unity ionization fraction (an upper bound), the overall contribution to the O VII column density is still small ($\lesssim 30\%$) compared to the value observed.

The sight line to 3C 273 extends into the Galactic halo through the edges of Radio Loop I and IV, which have been attributed to supernova remnants (see, e.g., Egger & Aschenbach 1995). X-ray observations in this direction reveal a hot, X-ray emitting gas with a temperature of $T \sim 3 \times 10^6$ K and emission measure ($n_e^2 d$) $\sim (1 - 2) \times 10^{-2}$ cm $^{-6}$ pc, where d is the distance through Loops I and IV (Iwan 1980). This gives a rough estimate of electron density $n_e \approx 0.01$ cm $^{-3}$. Using a path length of < 350 pc through Loops I and IV (Berkhuijsen 1973) and an O VII ionization fraction of 0.01 at $T \sim 3 \times 10^6$ K in collisional ionization equilibrium, the estimated column density for solar-abundance gas is $\lesssim 8 \times 10^{15}$ cm $^{-2}$, which could be as much as $\sim 50\%$ of the observed O VII column density if the loops really are this large. (Iwan 1980 estimates a size $\lesssim 100$ pc.) However, since the O VII ionization fraction raises quickly to 1 as the temperature approaches 10^6 K, a small uncertainty on T could cause a large variation in the predicted column density.

4. Summary

Our *Chandra* observations indicate that a large amount of hot gas exists in the Local Group and/or in the Milky Way. The O VII column density in this direction is a factor of 2-2.5 higher than toward PKS 2155-304 (Nicastro et al. 2002; Fang et al. 2002) and a factor of 2-5 higher than toward H 1821+643 (Mathur, Weinberg, & Chen 2002), which combined with the strong X-ray emission from the North Polar Spur in the direction of 3C 273, suggests that the absorption we see may well be from a combination of Galactic and Local Group sources. We constrain the temperature to $2.3 \times 10^5 < T < 1.2 \times 10^6$ K and find the baryon overdensity δ_b is ~ 100 , assuming a standard β model for the intragroup medium. The derived properties

are consistent with those of the WHIM gas which are predicted from simulations. For the case of Galactic absorption, we have outlined several scenarios to explain the observed O VII absorption, which may be associated with O VI observed at UV wavelengths. We find that in both the Galactic or Local Group cases there is a strong indication that collisional ionization is the dominant ionization source, at least for the gas which produces O VII absorption. We expect future X-ray observations may provide more information on the distribution of O VII at $z \sim 0$ and its strength in different directions. Such observations would provide valuable insight needed to determine the the location of the hot gas.

We obtained the data from the *Chandra* Data Archive built by the *Chandra* X-ray Center. We also thank the members of the MIT/CXC team for their support. This work is supported in part by contracts NAS 8-38249 and SAO SV1-61010. KRS acknowledges financial support through NASA contract NAS5-32985 and Long Term Space Astrophysics grant NAG5-3485.

[Note: Recently, we learned from a conference presentation that Rasmussen et al. (2002) reported the detection of a similar feature in the XMM spectrum of 3C 273, which confirms our detection.]

REFERENCES

- Aldcroft, T., Elvis, M., McDowell, J & Fiore, F. 94, ApJ, 437, 584
- Benjamin, R. A. & Shapiro, P. R. 2000, ApJS, submitted
- Berkhuijsen, E. M. 1973, A&A, 24, 143
- Cavaliere, A. & Fusco-Femiano, R. 1976, A&A, 49, 137
- Cen, R., Kang, H., Ostriker, J. P., & Ryu, D. 1995, ApJ, 451, 436
- Cen, R. & Ostriker, J. P. 1999, ApJ, 514, 1
- Chen, X., Weinberg, D. H., Katz, N. & Davé, R. 2002, ApJ, submitted (astro-ph/0203319)
- Courteau, S. & van den Bergh, S. 1999, AJ, 118, 337
- Davé, R. et al. 2001, ApJ, 552, 473
- Egger, R. J. & Aschenbach, B. 1995, A&A, 294, L25
- Fabian, A. C. & Barcons, X. 1992, ARA&A, 30, 429
- Fang, T. & Canizares, C. R. 2000, ApJ, 539, 532
- Fang, T., Marshall, H. L., Lee, J. C., Davis, D. S., & Canizares, C. R. 2002, ApJ, 572, L127
- Ferland, G. J., Korista, K. T., Verner, D. A., Ferguson, J. W., Kingdon, J. B., & Verner, E. M. 1998, PASP, 110, 761
- Fukugita, M., Hogan, C. J. & Peebles, J. P. E. 1998, ApJ, 503, 518
- Haardt, F. & Madau, P. 1996, ApJ, 461, 20
- Heckman, T. M., Norman, C. A., Strickland, D. K., & Sembach, K. R. 2002, ApJ, submitted (astro-ph/0205556)
- Hellsten, U., Gnedin, N. Y. & Miralda-Escudé, J. 1998, ApJ, 509, 56
- Houck, J. C. & Denicola, L. A. 2000, ASP Conf. Ser. 216: Astronomical Data Analysis Software and Systems IX, 9, 591
- Iwan, D. 1980, ApJ, 239, 316
- Kaspi, S. et al. 2002, ApJ, 574, 643

- Lockman, F. J. & Savage, B. D. 1995, ApJS, 97, 1
- Mathur, S., Weinberg, D., & Chen, X. 2002, ApJ, submitted (astro-ph/0206121)
- Miyaji, T., Ishisaki, Y., Ogasaka, Y., Ueda, Y., Freyberg, M. J., Hasinger, G., & Tanaka, Y. 1998, A&A, 334, L13
- Mulchaey, J. S., Davis, D. S., Mushotzky, R. F., & Burstein, D. 1996, ApJ, 456, 80
- Nicastro, F. et al. 2002, ApJ, 573, 157
- Nicastro, F. et al. 2002, Nature, submitted (astro-ph/0208012)
- Ostriker, J. P. & Cen, R. 1996, ApJ, 464, 27
- Perna, P. & Loeb, A. 1998, ApJ, 503, L135
- Rasmussen, J. & Pedersen, K. 2001, ApJ, 559, 892
- Rasmussen, A. et al. 2002, presentation in “the IGM/Galaxy Connection - The Distribution of Baryons at $z=0$ ”, Colorado
- Sanders, W.T., Edgar, R.J., Kraushaar, W. L., McCammon, D., & Morgenthaler, J.P. 2001, ApJ, 554, 694
- Sarazin, C. L. 1989, ApJ, 345, 12
- Savage, B. D. et al. 2000, ApJ, 538, L27
- Sembach, K. R. et al. 2000, ApJ, 538, L31
- Savage, B.D., et al. 2002, ApJS, submitted (astro-ph/0208140)
- Sembach, K. R. 2001, Extragalactic Gas at Low Redshift, eds. Mulchaey & Stocke, ASP Conference Series, in press (astro-ph/0108088)
- Sembach, K. R., Howk, J. C., Savage, B. D., Shull, J. M., & Oegerle, W. R. 2001, ApJ, 561, 573
- Sembach, K. R. et al. 2002, ApJS, submitted (astro-ph/0207562)
- Sfeir, D. M., Lallement, R., Crifo, F., & Welsh, B. Y. 1999, A&A, 346, 785
- Shapiro, P. R. & Bahcall, J. N. 1980, ApJ, 241, 1
- Shapiro, P. R. & Moore, R. T. 1976, ApJ, 207, 460

- Shull, J. M., Roberts, D., Giroux, M. L., Penton, S. V., & Fardal, M. A. 1999, *AJ*, 118, 1450
- Slavin, J. D., Shull, J. M., & Begelman, M. C. 1993, *ApJ*, 407, 83
- Snowden, S. L., Egger, R., Finkbeiner, D. P., Freyberg, M. J., & Plucinsky, P. P. 1998, *ApJ*, 493, 715
- Spitzer, L. 1978, *Physical Processes in the Interstellar Medium* (Princeton: Princeton University Press)
- Sutherland, R. S. & Dopita, M. A. 1993, *ApJS*, 88, 253
- Tripp, T. M., Savage, B. D., & Jenkins, E. B. 2000, *ApJ*, 534, L1
- Verner, D. A., Verner, E. M., & Ferland, G. J. 1996, *Atomic Data and Nuclear Data Tables*, 64, 1
- Wakker, B. P. et al. 2002, *ApJS*, submitted (astro-ph/0208009)

Table 1: Line Fitting Parameters

	O VII He α	O VII He β	O VIII Ly α
Wavelength ($\text{\AA}$)	21.60 ± 0.01	...	...
Rest Wavelength ($\text{\AA}$) ^a	21.6019	18.6288	18.9689
cz (km s ⁻¹)	-26 ± 140	...	...
Line Width ($\text{\AA}$) ^b	< 0.02	...	...
Line Flux ^c	$4.15^{+1.83}_{-0.90}$	...	...
W_λ (m $\text{\AA}$)	$28.4^{+12.5}_{-6.2}$	$< 11.9^d$	$< 14.2^d$
S/N	6.4	...	...

a. Verner, Verner, & Ferland (1996).

b. 90% upper limit of the line width σ .

c. Absorbed line flux, in units of 10^{-5} photons cm⁻²s⁻¹.

d. $4\text{-}\sigma$ detection upper limit.

Fig. 1.— The *Chandra* LETG-ACIS spectrum of 3C273 between 19 and 23 Å. The red solid line is the fitted continuum (Galactic absorbed power law plus the polynomial) plus the Gaussian model. The average 1σ error bar plotted on the right is based on statistics only. The inset shows the wavelength region where the rest-frame O VIII Ly α and O VII He β are located.

Fig. 2.— O VII column density versus central baryon density. The four different colors denote four different Local Group temperatures [$T = 2.3^5$ K (black), 5.8×10^5 K (red), 1.2×10^6 K (green) and 3.5×10^6 K (blue)]. The shadowed area indicates the 90% confidence range of the detected O VII column density. The two vertical dashed lines indicate the central density lower limit (from path length constraints) and upper limit (from $N_{\text{O VII}}$).

Fig. 3.— The column density of O VII versus O VI (solid line) and O VIII versus O VI (dashed line). The blue region indicates the allowed range from the measured $N(\text{O VII})/N(\text{O VI})$ ratio. The red line is the lower limit of the $N(\text{O VIII})/N(\text{O VI})$ from the non-detection of O VIII in the *Chandra* spectra.





